

FAX: 866-455-5150

**Supplier Information** 

VasoCARE, LLC 6554 Florida Blvd, Suite 123 Baton Rouge, LA 70806 PHONE: 800-256-9979

Fax 866-455-5150 NPI: 1720256662

## Prescription/Rx for ordering Comprehensive Pneumatic Compression, Neuromuscular Electrical Stimulation, or DVT Prevention Therapy Program

Progress notes must be attached and faxed along with this completed Rx form.

Patient's Name:	DOB:	Phone:
☐ Pneumatic Compression Therapy Program	<ul> <li>□ Neuromuscular Electrical Stimulation Therapy Program         Check the corresponding diagnosis:         <ul> <li>□ Disuse Muscle Atrophy (M62.561) Right Lower Limb</li> <li>□ Disuse Muscle Atrophy (M62.562) Left Lower Limb</li> <li>□ Disuse muscle atrophy due to lack of walking due to chronic disease or injury.</li> <li>□ Nerve supply is intact with the calf muscle.</li> </ul> </li> <li>Limb Pain or Ulcers due to PAD lead to disuse muscle atrophy:         <ul> <li>○ Claudication Limb Pain due to PAD</li> <li>○ Chronic edema due to Chronic Venous Insufficiency</li> <li>○ Diabetic Foot ulcer due to Arterial Insufficiency</li> <li>○ Restless Leg Syndrome (Movement Disorder due to CVI)</li> </ul> </li> </ul>	
Check the corresponding diagnosis:		
Lymphedema is Secondary to other Causes: I89.0		
Lymphedema Secondary to Post Mastectomy: 197.2		
_ Primary Lymphedema including Tarda: Q82.0		
Lymphedema due to Chronic Venous Insufficiency: 187.2		
CVI with Venous Stasis Ulcers {I87.331 Rt. or I87.332 Lt.}		
Patient compliant with instructions:		
$\square$ Elevate extremities daily to reduce swelling		
☐ Exercise daily to increase fluid flow and reduce swelling		
☐ Wear 30-mmHg compression garments for swelling		
Patient was last seen by physician after a 4-week trial		uctions: NMES Device - E-0745
of conservative therapies: DATE//	☐ Perform one hou	ur session times per day
Treatment Instructions: E0650/E0651 E0652 (calibrated pressure)	<ul><li>□ DVT Prevention 1</li><li>_ Deep Vein Thron</li></ul>	Therapy Program  nbosis/Pulmonary Emboli Prevention
☐ Apply to lower extremity: Right Left Bilateral	Treat	tment Instructions:
☐ Apply to upper extremity: Right Left Bilateral	☐ Wear DVT device	throughout day/evening hours to prevent
☐ Default Pressure Level: 40-60 mmHg, 60 minutes, BID	DVT and manage	post-surgical edema until fully ambulatory.
FACE-TO-FACE ENCOU	JNTER CERTIFICATION	I
I certify this patient is under my care, has been seen within 6 months of this order	er, supports the CMS LCD policy, a	and meets the need of this medical equipment.
Drocaribor Names	NDI.	Order Date.
Prescriber Name:PRINT	NPI:	Order Date:
Signature:		
Address/City/State/Zip:		
Email: Phone:		Fax:
Received by: Please fax: Rx order form, demogration	raphics sheet, and cl	inical notes to:
,	@ 866-455-5150	
	OR .	
	Myasocare com	